

Variability in Convective Feature Populations with ITCZ Width in the Pacific




Anita D. Rapp and Kyle Wodzicki

6 November 2019




"Deep Tropics Squeeze"





Lau and Kim 2015


"Tightening of Hadley Ascent"



Su et al. 2017


ITCZ Location, Width, Intensity Metrics

- Modified version of Berry and Reeder (2014) identification method
 - Apply a series of dynamic and thermodynamic masks to monthly mean ERA-Interim reanalysis
 - Divergence, ∇ Divergence, ∇ 2Divergence, $\theta_w(850)$





Use monthly mean rain rates to identify ITCZ boundaries

ITCZ Identification Example



Long-term Pacific ITCZ Trends





From Wodzicki and Rapp (2016)

Byrne et al. (2018)

Stephens et al. (2018)

Summary


- Observations show long-term narrowing of ITCZ, especially in central Pacific
- Precipitation intensity in ITCZ center increases with narrowing
- Dry layer frequency in ITCZ region has increased, especially in region of greatest narrowing (not shown)

Current work


How does the population of individual precipitating cloud systems vary with large-scale ITCZ variations?

Precipitation Features (PFs)

Extracted all TRMMPR PFs from Liu et al.(2008) that fall withinITCZ boundaries





Composited distribution of PF anomalies for widest and narrowest 25% of ITCZ width distribution




Liu et al. (2008)

PF Maximum Height






PF Area



PF Stratiform Area



Summary

- When ITCZ is wide:
 - PFs deeper and larger, with greater stratiform area
- When ITCZ is narrow:
 - PFs smaller and shallow

Future work

☼ Build a cloud feature dataset with VIRS, TMI, and PR to examine precipitation-anvil cloud area relationships change with environment, ENSO, large-scale ITCZ state?

